

Please write clearly in block capitals.

Centre number

Candidate number

Surname Lotfi-Baker  
Nada

Forename(s) www.onlinetutors.xyz

Candidate signature 07854144887

# A-level MATHEMATICS

Paper 3

Time allowed: 2 hours

## Materials

- You must have the AQA Formulae for A-level Mathematics booklet.
- You should have a graphical or scientific calculator that meets the requirements of the specification.

## Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

## Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 100.

## Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.


For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
<b>TOTAL</b>	



## Section A

Do not write  
outside the  
boxAnswer **all** questions in the spaces provided.

- 1 The graph of  $y = \arccos x$  is shown.

State the coordinates of the end point  $P$ .

Circle your answer.

**[1 mark]**

$(-\pi, 1)$

$(-1, \pi)$

$(-\frac{\pi}{2}, 1)$

$(-1, \frac{\pi}{2})$



2 Simplify fully

$$\frac{(x+3)(6-2x)}{(x-3)(3+x)} \quad \text{for } x \neq \pm 3$$

Circle your answer.

$$\textcircled{-2}$$

2

$$\frac{6-2x}{x-3}$$

$$\frac{2x-6}{x-3}$$

[1 mark]

3  $f(x) = 3x^2$

Obtain  $\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

Circle your answer.

$$\frac{3h^2}{h}$$

 $x^3$ 

$$\frac{3(x+h)^2 - 3x^2}{h}$$

$$\textcircled{6x}$$

[1 mark]

Turn over for the next question

Turn over ►



- 4 (a) Show that the first three terms, in descending powers of  $x$ , of the expansion of

$$(2x - 3)^{10}$$

are given by

$$1024x^{10} + px^9 + qx^8$$

where  $p$  and  $q$  are integers to be found.

[3 marks]

$$\begin{aligned} & (2x)^{10} + {}^{10}C_1(2x)^9(-3) + {}^{10}C_2(2x)^8(-3)^2 \\ & = 1024x^{10} - 15360x^9 + 103680x^8 \end{aligned}$$

- 4 (b) Find the constant term in the expansion of


$$\left(2x - \frac{3}{x}\right)^{10}$$

[2 marks]

$${}^{10}C_5 \times 2^5 \times (-3)^5 = 1959552$$



- 5 A gardener is creating flowerbeds in the shape of sectors of circles.
- The gardener uses an edging strip around the perimeter of each of the flowerbeds.
- The cost of the edging strip is £1.80 per metre and can be purchased for any length.
- One of the flowerbeds has a radius of 5 metres and an angle at the centre of 0.7 radians as shown in the diagram below.



- 5 (a) (i) Find the area of this flowerbed.

[2 marks]

$$\begin{aligned} \text{Area of a sector} &= \frac{1}{2} \times 5^2 \times 0.7 \\ \text{(in radians } \frac{1}{2} r^2 \theta) & \\ &= 8.75 \text{ m}^2 \end{aligned}$$


---



---



---



---

Question 5 continues on the next page

Turn over ►



5 (a) (ii) Find the cost of the edging strip required for this flowerbed.

[3 marks]

$$\begin{aligned} \text{arc length} &= r\theta \\ &= 0.7(5) = 3.5 \end{aligned}$$

$$\begin{aligned} \text{total perimeter} &= 3.5 + 5 + 5 \\ &= 13.5\text{m} \end{aligned}$$

$$\begin{aligned} \text{cost} &: 13.5\text{m} \times \pounds 1.80 \\ &= \pounds 24.30 \end{aligned}$$

5 (b) A flowerbed is to be made with an area of  $20\text{m}^2$

5 (b) (i) Show that the cost,  $\pounds C$ , of the edging strip required for this flowerbed is given by

$$C = \frac{18}{5} \left( \frac{20}{r} + r \right)$$

where  $r$  is the radius measured in metres.

[3 marks]

$$\text{perimeter: } p = r\theta + 2r$$

$$\text{area: } \frac{1}{2} r^2 \theta = 20 \rightarrow \theta = \frac{40}{r^2}$$

$$p = r \left( \frac{40}{r^2} \right) + 2r$$

$$p = \frac{40}{r} + 2r$$

$$C = \frac{40 \times 1.8}{r} + 2 \times 1.8 r$$

$$= \frac{72}{r} + \frac{18}{5} r = \frac{18}{5} \left( \frac{20}{r} + r \right)$$



- 5 (b) (ii) Hence, show that the minimum cost of the edging strip for this flowerbed occurs when  $r \approx 4.5$

Fully justify your answer.

[5 marks]

we know  $C = \frac{72}{r} + \frac{18}{5}r$

$$\frac{dC}{dr} = -\frac{72}{r^2} + \frac{18}{5}$$

minimum occurs when  $\frac{dC}{dr} = 0$

$$-\frac{72}{r^2} + \frac{18}{5} = 0 \quad r^2 = 20$$

$$r = \sqrt{20} \approx 4.472$$

$$r \approx 4.5$$

check this is a minimum:

$$\frac{d^2C}{dr^2} = \frac{144}{r^3} \quad \text{when } r = \sqrt{20} \quad \frac{d^2C}{dr^2} > 0$$

therefore minimum at  $r \approx 4.5$



- 6 Given that  $x > 0$  and  $x \neq 25$ , fully simplify

$$\frac{10 + 5x - 2x^{\frac{1}{2}} - x^{\frac{3}{2}}}{5 - \sqrt{x}}$$

Fully justify your answer.

[4 marks]

$$\begin{aligned} & \frac{10 + 5x - 2x^{1/2} - x^{3/2}}{5 - \sqrt{x}} \times \frac{5 + \sqrt{x}}{5 + \sqrt{x}} \\ = & \frac{\cancel{50} + 10\sqrt{x} + \cancel{25x} + 5x\sqrt{x} - \cancel{10\sqrt{x}} - \cancel{2x} - \cancel{5x^{3/2}} - \cancel{x^{3/2}}\sqrt{x}}{25 - x} \\ = & \frac{\cancel{50} + 23x - x^2}{25 - x} \\ = & \frac{(25 - x)(2 + x)}{25 - x} = 2 + x \end{aligned}$$

Do not write  
outside the  
box



**Turn over for the next question**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**

**Turn over ▶**



7 A building has a leaking roof and, while it is raining, water drips into a 12 litre bucket.

When the rain stops, the bucket is one third full.

Water continues to drip into the bucket from a puddle on the roof.

In the first minute after the rain stops, 30 millilitres of water drips into the bucket.

In each subsequent minute, the amount of water that drips into the bucket reduces by 2%.

During the  $n$ th minute after the rain stops, the volume of water that drips into the bucket is  $W_n$  millilitres.

7 (a) Find  $W_2$

[1 mark]

$$30 \times 0.98 = 29.4$$

$$W_2 = 29.4$$

7 (b) Explain why

$$W_n = A \times 0.98^{n-1}$$

and state the value of  $A$ .

[2 marks]

$W_n$  is the  $n$ th term of a geometric sequence,

a 2% reduction gives a common ratio of 0.98.

$$A = 30$$



- 7 (c) Find the increase in the water in the bucket 15 minutes after the rain stops.

Give your answer to the nearest millilitre.

[2 marks]

$$S_n = \frac{a(1-r^n)}{(1-r)}$$

$$S_{15} = \frac{30(1-0.98^{15})}{1-0.98} = 392$$

- 7 (d) Assuming it does not start to rain again, find the maximum amount of water in the bucket.

[3 marks]

$$S_{\infty} = \frac{a}{1-r}$$

$$= \frac{30}{1-0.98} = 1500 = 1.5 \text{ litres}$$

$$1.5 + 4 = 5.5 \text{ litres}$$

- 7 (e) After several hours the water has stopped dripping.

Give **two** reasons why the amount of water in the bucket is not as much as the answer found in part (d).

[2 marks]

° Sum to infinity was used but this assumes there are infinite drips, but they have stopped

° Water will evaporate over time

Turn over ►



8

Given that

$$\int_{\frac{\pi}{4}}^{\frac{\pi}{3}} x \cos x \, dx = a\pi + b$$

find the exact value of  $a$  and the exact value of  $b$ .

Fully justify your answer.

**[6 marks]**

integration by parts:

$$u = x$$

$$u' = 1$$

$$v' = \cos x$$

$$v = \sin x$$

$$\int x \cos x \, dx = x \sin x - \int \sin x \, dx$$

$$= x \sin x + \cos x$$

$$\int_{\frac{\pi}{4}}^{\frac{\pi}{3}} x \cos x \, dx = [x \sin x + \cos x]_{\frac{\pi}{4}}^{\frac{\pi}{3}}$$

$$= \frac{\pi}{3} \sin \frac{\pi}{3} + \cos \frac{\pi}{3} - \left( \frac{\pi}{4} \sin \frac{\pi}{4} + \cos \frac{\pi}{4} \right)$$

$$= \pi \frac{\sqrt{3}}{6} + \frac{1}{2} - \left( \pi \frac{\sqrt{2}}{8} + \frac{\sqrt{2}}{2} \right)$$

$$= \left( \frac{4\sqrt{3} - 3\sqrt{2}}{24} \right) \pi + \left( \frac{1 + \sqrt{2}}{2} \right)$$

Do not write  
outside the  
box



- 9 A function  $f$  is defined for all real values of  $x$  as

$$f(x) = x^4 + 5x^3$$

The function has exactly two stationary points when  $x = 0$  and  $x = -\frac{15}{4}$

- 9 (a) (i) Find  $f''(x)$

[2 marks]

$$f(x) = x^4 + 5x^3$$

$$f'(x) = 4x^3 + 15x^2$$

$$f''(x) = 12x^2 + 30x$$

- 9 (a) (ii) Determine the nature of the stationary points.

Fully justify your answer.

[4 marks]

$$\text{at } x=0 \quad f''(x) = 0$$

$$\text{check } f''(1) = 12 + 30 > 0$$

$$f''(-1) = 12 - 30 < 0$$

$\therefore x=0$  is a point of inflection

$$\text{at } x = -\frac{15}{4} \quad f''(x) = 12\left(-\frac{15}{4}\right)^2 + 30\left(-\frac{15}{4}\right)$$

$$= \frac{225}{4} > 0$$

$\therefore x = -\frac{15}{4}$  is a minimum point



- 9 (b) State the range of values of  $x$  for which

$$f(x) = x^4 + 5x^3$$

is an increasing function.

[1 mark]

$$x > -15/4$$

- 9 (c) A second function  $g$  is defined for all real values of  $x$  as

$$g(x) = x^4 - 5x^3$$

- 9 (c) (i) State the single transformation which maps  $f$  onto  $g$ .

[1 mark]

Reflection in the  $y$  axis

- 9 (c) (ii) State the range of values of  $x$  for which  $g$  is an increasing function.

[1 mark]

$$x > 15/4$$

Turn over ►



## Section B

Answer **all** questions in the spaces provided.

- 10 Anke has collected data from 30 similar-sized cars to investigate any correlation between the age of the car and the current market value.

She calculates the correlation coefficient.

Which of the following statements best describes her answer of  $-1.2$ ?

Tick (✓) **one** box.

[1 mark]

Definitely incorrect

Probably incorrect

Probably correct

Definitely correct

- 11 The random variable  $X$  is such that  $X \sim B(n, p)$

The mean value of  $X$  is 225

The variance of  $X$  is 144

Find  $p$ .

Circle your answer.

[1 mark]

0.36

0.6

0.64

0.8



12

An electoral register contains 8000 names.

A researcher decides to select a systematic sample of 100 names from the register.

Explain how the researcher should select such a sample.

**[3 marks]**

Give each name a number from 1 to 8000

$$\frac{8000}{100} = 80$$

Randomly select a number between 1 and 80 and select every 80th person from the first person selected

Turn over for the next question

Turn over ►



13 The table below is an extract from the Large Data Set.

Propulsion Type	Region	Engine Size	Mass	CO <sub>2</sub>	Particulate Emissions
2	London	1896	1533	154	0.04
2	North West	1896	1423	146	0.029
2	North West	1896	1353	138	0.025
2	South West	1998	1547	159	0.026
2	London	1896	1388	138	0.025
2	South West	1896	1214	130	0.011
2	South West	1896	1480	146	0.029
2	South West	1896	1413	146	0.024
2	South West	2496	1695	192	0.034
2	South West	1422	1251	122	0.025
2	South West	1995	2075	175	0.034
2	London	1896	1285	140	0.036
2	North West	1896	0	146	

13 (a) (i) Calculate the mean and standard deviation of CO<sub>2</sub> emissions in the table.

[2 marks]

$$\text{Mean} = 148.6$$

$$\text{Standard deviation} = 17.8$$



- 13 (a) (ii) Any value more than 2 standard deviations from the mean can be identified as an outlier.

Determine, using this definition of an outlier, if there are any outliers in this sample of CO<sub>2</sub> emissions.

Fully justify your answer.

[2 marks]

$$148.6 + 2(17.8) = 184.2$$

$$148.6 - 2(17.8) = 113$$

$$113 < \text{outlier} \quad 184.2 < \text{outlier}$$

$$192 > 184.2$$

$\therefore$  192 is the only outlier

- 13 (b) Maria claims that the last line in the table must contain two errors.

Use your knowledge of the Large Data Set to comment on Maria's claim.

[2 marks]

• The 0 value is an error because every car has a mass

• The blank cell may not be an error as not all particulate emissions are recorded in the LDS

Turn over for the next question

Turn over ►



14  $A$  and  $B$  are two events such that

$$\text{both} \rightarrow P(A \cap B) = 0.1$$

$$\text{not } A \text{ + not } B \rightarrow P(A' \cap B') = 0.2$$

$$P(B) = 2P(A)$$

14 (a) Find  $P(A)$

[4 marks]

~~the answer~~

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A \cup B) = 0.8$$

$$P(A \cup B) = P(A) + 2P(A) - 0.1$$

$$0.9 = 3P(A)$$

$$P(A) = 0.3$$

14 (b) Find  $P(B|A)$

[2 marks]

$$P(A \cap B) = P(A) \times P(B|A)$$

$$P(B|A) = \frac{P(A \cap B)}{P(A)}$$

$$= \frac{0.1}{0.3} = \frac{1}{3}$$



14 (c) Determine if  $A$  and  $B$  are independent events.

[1 mark]

not independent as  $P(A) \times P(B) = 0.3 \times 0.6 =$   
 $0.18 \neq P(A \cap B) = 0.1$

---

Turn over for the next question

Turn over ►



15

A team game involves solving puzzles to escape from a room.

Using data from the past, the mean time to solve the puzzles and escape from one of these rooms is 65 minutes with a standard deviation of 11.3 minutes.

After recent changes to the puzzles in the room, it is claimed that the mean time to solve the puzzles and escape has changed.

To test this claim, a random sample of 100 teams is selected.

The total time to solve the puzzles and escape for the 100 teams is 6780 minutes.

Assuming that the times are normally distributed, test at the 2% level the claim that the mean time has changed.

[7 marks]

$X$  = time to solve in minutes

$$H_0: \mu = 65$$

$$H_1: \mu \neq 65$$

$$\begin{aligned} \text{Test statistic} &= \frac{67.8 - 65}{(11.3 / \sqrt{100})} \\ &= 2.48 \end{aligned}$$

$$\text{Critical value} = 2.33$$

$$2.48 > 2.33$$

$\therefore$  Reject  $H_0$ : there is sufficient evidence at the 2% level to suggest that mean escape time has changed.



16

The discrete random variable  $X$  has the probability function

$$P(X=x) = \begin{cases} c(7-2x) & x=0, 1, 2, 3 \\ k & x=4 \\ 0 & \text{otherwise} \end{cases}$$

where  $c$  and  $k$  are constants.

16 (a)

Show that  $16c + k = 1$ 

[2 marks]

$x$	0	1	2	3	4
$P(X=x)$	$7c$	$5c$	$3c$	$c$	$k$

$$7c + 5c + 3c + c + k = 1$$

$$16c + k = 1$$

16 (b)

Given that  $P(X \geq 3) = \frac{5}{8}$ find the value of  $c$  and the value of  $k$ .

[2 marks]

$$c + k = \frac{5}{8}$$

$$c = \frac{1}{40} \quad k = \frac{3}{5}$$

Turn over for the next question

Turn over ►



17 James is playing a mathematical game on his computer.

The probability that he wins is 0.6

As part of an online tournament, James plays the game 10 times.

Let  $Y$  be the number of games that James wins.

17 (a) State two assumptions, in context, for  $Y$  to be modelled as  $B(10, 0.6)$

[2 marks]

• The event of James winning one game is independent of him winning another game

• The probability of James winning remains constant at 0.6 from game to game.

17 (b) Find  $P(Y = 4)$

[1 mark]

$$0.111$$

17 (c) Find  $P(Y \geq 4)$

[2 marks]

$$P(Y \leq 3) = 0.05476$$

$$P(Y \geq 4) = 1 - P(Y \leq 3)$$

$$= 1 - 0.05476$$

$$= 0.94524$$



- 17 (d) After practising the game, James claims that he has increased his probability of winning the game.

In a random sample of 15 subsequent games, he wins 12 of them.

Test a 5% significance level whether James's claim is correct.

[6 marks]

$X =$  number of games won

$$H_0: p = 0.6$$

$$H_1: p > 0.6$$

$$X \sim B(15, 0.6)$$

$$P(X \geq 12) = 1 - P(X \leq 11)$$

$$= 1 - 0.9094$$

$$= 0.0905$$

$$0.0905 > 0.05 \text{ so accept } H_0$$

There is insufficient evidence to suggest that the probability of James winning the game has increased

Turn over for the next question

Turn over ►



18 A factory produces jars of jam and jars of marmalade.

18 (a) The weight,  $X$  grams, of jam in a jar can be modelled as a normal variable with mean 372 and a standard deviation of 3.5

18 (a) (i) Find the probability that the weight of jam in a jar is equal to 372 grams.

[1 mark]

0

18 (a) (ii) Find the probability that the weight of jam in a jar is greater than 368 grams.

[2 marks]

$$P(X > 368) = 0.87345$$

18 (b) The weight,  $Y$  grams, of marmalade in a jar can be modelled as a normal variable with mean  $\mu$  and standard deviation  $\sigma$

18 (b) (i) Given that  $P(Y < 346) = 0.975$ , show that

$$346 - \mu = 1.96\sigma$$

Fully justify your answer.

[3 marks]

Using inverse normal, z value is 1.95996...

for the area of 0.975

$$P\left(Z < \frac{346 - \mu}{\sigma}\right) = 0.975$$

$$\frac{346 - \mu}{\sigma} = 1.96 \quad \therefore 346 - \mu = 1.96\sigma$$



18 (b) (ii) Given further that

$$P(Y < 336) = 0.14$$

find  $\mu$  and  $\sigma$

[4 marks]

using inverse normal distribution again:

$$z = -1.08$$

$$\frac{336 - \mu}{\sigma} = -1.08$$

$$336 - \mu = -1.08\sigma$$

$$\sigma = 3.29$$

$$\mu = 340$$

END OF QUESTIONS



**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**









